

Shawn Hennigan, MD Surgery Pain Medication Protocol

David A. Toivonen, M.D.

Jon J. Cherney, M.D.

Boyd C. Lumsden, M.D.

Scott P. Olvey, M.D.

Joseph P. Cullen, M.D.

Nathan L. Van Zeeland, M.D.

Matthew A. Butler, M.D.

Shawn P. Hennigan, M.D.

Blake W. Hildahl, M.D., ATC

Administrator:

Tina M. Sauer, CAPPM

Prior to surgery:

- 1. Start your *Mobic* prescription the night prior to surgery
 - Take 1 full tablet the night prior to surgery. Starting the day of surgery,
 take a ½ tablet twice daily (AM and PM). Take this until your 1st follow up
 appointment, and then as needed.

It is important to avoid taking other anti-inflammatory medications such as Advil, Ibuprofen, Aleve, Naproxen, Celebrex, ect, while on this prescription

- 2. Start your *gabapentin* prescription 1 week prior to surgery.
 - Take 1 tablet at nighttime starting 1 week prior to surgery. Starting the day of surgery, increase to 1 tablet twice daily (AM and PM). Take this until your 1st follow up appointment.

After Surgery:

Post-op Days 1-2:

- 1. *Oxycodone 5mg*: Take 1-2 tablets every 4-6 hours as needed for pain.
- 2. *Mobic*: Continue a ½ tablet twice daily.
- 3. *Gabapentin:* Continue 1 tablet twice daily
- **4.** Extra Strength Tylenol/Acetaminophen 500mg tablets: You may take 2 tablets every 6 hours as needed, not to exceed 4,000mg per day.

Post-op Days 3-10:

- 1. *Oxycodone 5mg*: Decrease to 1-2 tablets at nighttime only until prescription is gone
- 2. *Mobic*: Continue a ½ tablet twice daily.
- 3. *Gabapentin:* Continue 1 tablet twice daily
- 4. Continue *Extra Strength Tylenol/Acetaminophen:* 2 tablets every 6 hours as needed, a max of 4,000mg per day.

2323 N. Casaloma Drive Appleton, WI 54913 P. 920-730-8833 TF. 888-383-3039 F. 920-738-1974

www.handtoshoulderwisconsin.com

Satellite Office Location: Green Bay

^{**}All above medications are safe to take simultaneously **